

# 1-Cohomology of Chevalley Groups

HELMUT VÖLKLEIN

*Department of Mathematics, University of Florida,  
Gainesville, Florida 32611*

*Communicated by Wilberd van der Kallen*

Received March 14, 1988; revised August 8, 1988

We relate the 1-cohomology of a finite Chevalley group to induced modules and finite geometry. As a corollary, we get the 1-cohomology of most adjoint modules. © 1989 Academic Press, Inc.

## INTRODUCTION

Let  $\Sigma$  be an irreducible root system and  $G$  the universal Chevalley group of type  $\Sigma$  over the field  $\mathbf{F}_q$  of  $q$  elements. In this paper we want to study the 1-cohomology  $H^1(G, W)$ , where  $W$  is an irreducible module for  $G$  over the algebraic closure  $K$  of  $\mathbf{F}_q$ . Let  $V := W^*$  be the dual of  $W$ , and  $B$  a Borel subgroup of  $G$ . By [CPSK] we know that if  $q$  is “large” (depending on  $\Sigma$  and  $W$ ) then any non-splitting extension

$$0 \rightarrow K \rightarrow E \rightarrow V \rightarrow 0$$

of  $KG$ -modules (where  $K$  is considered as trivial module) satisfies the “highest weight condition”:

$E$  is generated (as a  $KG$ -module) by a 1-dimensional subspace fixed by  $B$ . (\*)

By Frobenius reciprocity, the condition  $(*)$  is equivalent to the one that  $E$  is a quotient of the module  $\lambda_B^G$  induced from a 1-dimensional representation  $\lambda$  of  $B$ . So if for given  $G$  and  $V$ ,  $(*)$  holds for all extensions  $E$  of  $V$  over  $K$ , then  $H^1(G, W)$  can be read off from the induced module  $\lambda_B^G$ . (Thereby, we could replace  $\lambda_B^G$  by the (possibly) smaller module  $\lambda_P^G$ , where  $P$  is the full stabilizer in  $G$  of the  $B$ -invariant 1-space in  $V$ ; namely,  $(*)$  is equivalent to the same condition with  $B$  replaced by  $P$ .)

The main purpose of this paper is to verify  $(*)$  in many cases not covered by the stability results of [CPSK]. In fact much of the work is devoted to the case  $q \leq 3$ . In Section 2, we study the case that  $V$  is a fundamental module, and more generally, a module whose highest weight is a multiple

of a fundamental weight. In this case we see rather quickly that  $(*)$  holds for all  $q > 3$ . We also settle the case  $q \leq 3$  for the fundamental modules, by reducing to some explicit results on 1-cohomology from [JP]. (We need two more such computations, which we perform in the Appendix). In Section 3, we obtain some results in the case  $q > 2$ ; in particular, we prove that  $(*)$  always holds if  $q$  is not a prime. The idea is to reduce to the case  $G = SL_2(q)$ , where all projective indecomposable modules are known by [AJL].

Another consequence of  $(*)$  (which provided the original motivation for this paper), is the following: For each parabolic subgroup  $P$  of  $G$ , let  $V_P$  (resp.  $E_P$ ) denote the fixed space in  $V$  (resp. in  $E$ ) of the unipotent radical of  $P$ . By [S-1],  $V_P$  is an irreducible module for each Levi-complement  $L_P$  of  $P$ . Thus  $(*)$  implies that for each  $P$  we get an extension

$$0 \rightarrow K \rightarrow E_P \rightarrow V_P \rightarrow 0 \quad (+)$$

of  $L_P$ -modules. We are particularly interested in the case that  $P$  is a minimal parabolic. Then the "semisimple part" of  $L_P$  is isomorphic to  $SL_2(q)$ , and we can again apply the results of [AJL] to deduce splitting of  $(+)$  in many cases. If  $(+)$  splits for all minimal parabolics, then we can choose complements  $\bar{V}_P$  of  $K$  in  $E_P$  for all minimal parabolics  $P$ , and also for all Borel subgroups  $P$ , such that these spaces  $\bar{V}_P$  form the same incidence geometry in  $E$  as the corresponding spaces  $V_P$  in  $V$ . Thus  $E$  is generated (as a vector space) by copies of the spaces  $V_P$  (for  $P$  a minimal parabolic or Borel subgroup of  $G$ ), realising their incidence geometry from  $V$ ; and so  $E$  is a quotient of the "universal" module  $\tilde{V}$  so generated. (This module  $\tilde{V}$  was introduced in [RS-2, Section 3B] in a slightly different set-up.)

So under the above hypotheses,  $H^1(G, W)$  can be read off not only from  $\lambda_B^G$ , but already from its much smaller quotient  $\tilde{V}$ . Now  $\tilde{V}$  has been calculated for the "minimal" modules in [RS-1] and for the adjoint modules (not of type  $C_l$ ) in [V-1]. So as a corollary we get the 1-cohomology groups of these modules. Most of them have already been computed in [CPS], [JP] and [V-2], except for the adjoint modules of type  $B_l$ ,  $G_2$ , and  $F_4$  over fields with  $\leq 9$  elements.

This paper was partly motivated by [S-2], where the idea of studying the extension theory of irreducible  $KG$ -modules via the methods of [RS-1] is introduced.

## 1. NOTATIONS

The following notations will be kept fixed throughout the paper:

$\Sigma$  ... an irreducible root system in the Euclidean space  $E'$  with scalar product  $(\ , \ )$

- $\langle \ , \ \rangle \dots$  defined by  $\langle \alpha, \beta \rangle = 2(\alpha, \beta)/(\beta, \beta)$  for  $\alpha, \beta \in E'$   
 $r_\alpha \dots$  the reflection at  $\alpha$ , defined by  $r_\alpha(\beta) = \beta - \langle \beta, \alpha \rangle \alpha$   
 $\Delta = \{\alpha_1, \dots, \alpha_l\} \dots$  a system of simple roots, defining as usual a partial order on  $\Sigma$   
 $\Sigma^+ \dots$  the set of positive roots  
 $\lambda_1, \dots, \lambda_l \dots$  the corresponding fundamental weights, defined by  $\langle \lambda_i, \alpha_j \rangle = \delta_{i,j}$   
 $X \dots$  the weight lattice, spanned by  $\lambda_1, \dots, \lambda_l$   
 $q = p^n \dots$  a power of the prime  $p$   
 $k = \mathbb{F}_q \dots$  the field with  $q$  elements  
 $k^* \dots$  the multiplicative group of  $k$   
 $G \dots$  the universal Chevalley group of type  $\Sigma$  over  $k$ , constructed as in [St].

Then  $G$  comes with

- $h_\alpha: k^* \rightarrow G \dots$  coroot corresponding to  $\alpha \in \Sigma$   
 $T \dots$  Cartan subgroup generated by the  $h_\alpha(s)$ ,  $s \in k^*$ ,  $\alpha \in \Sigma$   
 $U_\alpha$  ( $\alpha \in \Sigma$ )  $\dots$  root subgroups  
 $B = TU \dots$  Borel subgroup, with  $U = O_p(B) = \prod_{\alpha > 0} U_\alpha$   
 $x_\alpha: k \rightarrow U_\alpha \dots$  additive isomorphisms satisfying Chevalley's commutator formula (see [St, p. 30, R2]) and the relations  $tx_\alpha(a)t^{-1} = x_\alpha(\alpha(t)a)$  for  $t \in T$ ,  $\alpha \in k$ . Thereby each  $\mu \in X$  acts on  $T$  via  $\mu(h_\alpha(s)) = s^{\langle \mu, \alpha \rangle}$ .

For each subset  $J$  of  $\Pi := \{1, \dots, l\}$  we set  $\Sigma_J := \Sigma \cap \langle \alpha_j: j \in J \rangle$  and  $G_J := \langle U_\alpha: \alpha \in \Sigma_J \rangle$ . If  $J$  is non-empty then  $\Sigma_J$  can naturally be considered as the root system of the Chevalley group  $G_J$ , and  $\Delta_J := \{\alpha_j: j \in J\}$  is a basis of  $\Sigma_J$ . Set  $L_J := G_J T$  and  $U_J := \langle U_\alpha: \alpha \in \Sigma^+ \setminus \Sigma_J \rangle$ . Then  $L_J$  (resp.  $U_J$ ) is a Levi complement (resp. the unipotent radical) of the parabolic subgroup  $P_J = L_J U_J$  of  $G$ .

Finally, we let  $A_i(q)$ ,  $B_i(q)$  etc. denote the *universal* Chevalley group of the corresponding type over  $k$ . The commutator subgroup of any group  $X$  is denoted  $X'$ , and  $C_Y(X)$  is the centralizer of  $X$  in  $Y$ .

## 2. MODULES WITH HIGHEST WEIGHT $m\lambda_j$

In this section,  $l = \text{rank } G > 1$ . We assume that for each irreducible root system  $\Sigma$  the simple roots  $\alpha_1, \dots, \alpha_l$  are labelled as in [Bou, Planches] except for type  $G_2$ , where we let  $\alpha_1$  be long. To avoid ambiguities in this labelling, we let type  $C_l$  (resp.  $D_l$ ) exist only for  $l > 2$  (resp.  $l > 3$ ).

LEMMA 1. Let  $\alpha \in \Sigma^+ \setminus \mathcal{A}$ . Then  $U_\alpha \subset U'$  unless

- (i)  $q = 2$ ,  $\Sigma \cong B_l$ , and  $\alpha = \alpha_{l-1} + \alpha_l$  or  $\alpha_{l-1} + 2\alpha_l$
- (i)'  $q = 2$ ,  $\Sigma \cong C_l$ , and  $\alpha = \alpha_{l-1} + \alpha_l$  or  $2\alpha_{l-1} + \alpha_l$
- (ii)  $q = 2$ ,  $\Sigma \cong F_4$ , and  $\alpha = \alpha_2 + \alpha_3$  or  $\alpha_2 + 2\alpha_3$
- (iii)  $q = 2$ ,  $\Sigma \cong G_2$ , and  $\alpha$  is short, or
- (iv)  $q = 3$ ,  $\Sigma \cong G_2$ , and  $\alpha = \alpha_1 + \alpha_2$  or  $\alpha_1 + 3\alpha_2$ .

*Proof.* We explicitly need the constants appearing in Chevalley's commutator formula (see [St, Lemma 33 and (3), p. 151]). Let  $\gamma, \delta \in \Sigma$  with  $|\gamma| \geq |\delta|$ . Define the commutator  $[g, h] = ghg^{-1}h^{-1}$  for  $g, h \in G$ , and let  $c, d \in k$ .

(a) If  $\gamma$  and  $\delta$  form a simple system of type  $A_2$ , then  $[x_\gamma(c), x_\delta(d)] = x_{\gamma+\delta}(cd)$ ; hence  $[U_\gamma, U_\delta] = U_{\gamma+\delta}$ .

(b) If  $\gamma$  and  $\delta$  form a simple system of type  $B_2$ , then

$$[x_\gamma(c), x_\delta(d)] = x_{\gamma+\delta}(\pm cd)x_{\gamma+2\delta}(\pm cd^2).$$

One checks easily that this implies  $[U_\gamma, U_\delta] = U_{\gamma+\delta}U_{\gamma+2\delta}$  if  $q \neq 2$ .

(c) If  $\gamma$  and  $\delta$  form a simple system of type  $G_2$ , then

$$[x_\gamma(c), x_\delta(d)] = x_{\gamma+\delta}(\pm cd)x_{\gamma+2\delta}(\pm cd^2)x_{\gamma+3\delta}(\pm cd^3)x_{2\gamma+3\delta}(\pm c^2d^3),$$

$$[x_{\gamma+3\delta}(c), x_\gamma(d)] = x_{2\gamma+3\delta}(\pm cd)$$

$$[x_{\gamma+\delta}(c), x_\delta(d)] = x_{\gamma+2\delta}(\pm 2cd)x_{\gamma+3\delta}(\pm 3cd^2)x_{2\gamma+3\delta}(\pm 3c^2d).$$

Case 1.  $\Sigma \not\cong G_2$ , and  $q \neq 2$  if  $\Sigma$  has two different root lengths.

Each  $\alpha \in \Sigma^+ \setminus \mathcal{A}$  can be written as  $\alpha = \alpha' + \alpha''$  with  $\alpha', \alpha'' \in \Sigma^+$ . If  $\alpha, \alpha'$ , and  $\alpha''$  have the same length, then  $\alpha'$  and  $\alpha''$  form a simple system of type  $A_2$ ; so  $U_\alpha = [U_{\alpha'}, U_{\alpha''}] \leq U'$ . Thus it remains to consider the case that  $q \neq 2$ , and  $\alpha', \alpha''$  span a subsystem of  $\Sigma$  of type  $B_2$ . If  $\alpha$  is short (resp. long), then  $\alpha'$  and  $\alpha''$  (esp.  $\alpha'$  and  $\alpha'' - \alpha'$ ) form a simple system of type  $B_2$ , hence  $U_\alpha \leq [U_{\alpha'}, U_{\alpha''}] \leq U'$  (resp.  $U_\alpha \leq [U_{\alpha'}, U_{\alpha'' - \alpha'}]$ ) by (b). If  $\alpha$  is long, note that we may assume that  $\alpha'' - \alpha'$  is positive (interchanging  $\alpha'$  and  $\alpha''$  if necessary).

Case 2.  $q = 2$  and  $\Sigma \cong B_l$ .

If  $\alpha \in \Sigma^+ \setminus \mathcal{A}$  is long and  $\neq \alpha_{l-1} + 2\alpha_l$ , then  $\alpha = \alpha' + \alpha''$  for long  $\alpha', \alpha'' \in \Sigma^+$  (see [Bou, Planche II]); so  $U_\alpha = [U_{\alpha'}, U_{\alpha''}] \leq U'$  as before. Now assume  $\alpha$  is short and  $\neq \alpha_{l-1} + \alpha_l$ . Then we can find long  $\gamma \in \mathcal{A}$  such that  $\delta := \alpha - \gamma \in \Sigma^+$  [Bou, Planche II]. Then  $\gamma$  and  $\delta$  form a simple system of type  $B_2$ , hence  $x_\alpha(c) = [x_\gamma(\pm c), x_\delta(1)]x_{\gamma+2\delta}(\pm c)$  by (b). So it suffices to show  $U_{\gamma+2\delta} \leq U'$ . Now  $\gamma + 2\delta \neq \alpha_{l-1} + 2\alpha_l$  (since otherwise  $\alpha = \gamma + \delta =$

$(\gamma + 2\delta) - \delta = \alpha_{l-1} + 2\alpha_l - \delta = \alpha_{l-1} + \alpha_l$  or  $= \alpha_l$ , which is excluded), and so  $U_{\gamma+2\delta} \leq U'$  by the above (since  $\gamma + 2\delta$  is long).

*Case 2'.  $q = 2$  and  $\Sigma \cong C_l$ .*

This reduces to Case 2 by the exceptional isomorphism  $B_l(2) \rightarrow C_l(2)$ .

*Case 3.  $q = 2$  and  $\Sigma \cong F_4$ .*

If  $\alpha \in \Sigma^+ \setminus \Delta$  is long and  $\neq \alpha_2 + 2\alpha_3$ ,  $\alpha_2 + 2\alpha_3 + 2\alpha_4$ , then  $\alpha = \alpha' + \alpha''$  for long  $\alpha'$ ,  $\alpha'' \in \Sigma^+$  (see [Bou, Planche VIII]); so  $U_\alpha \leq U'$  as before. Since  $\beta := \alpha_2 + 2\alpha_3 + 2\alpha_4 \in \Sigma_{\{2,3,4\}} \cong C_3$ , we have also  $U_\beta \leq U'$  by Case 2'. Finally we use the exceptional graph automorphism of  $F_4(2)$  (which interchanges long and short positive roots) to conclude that also  $U_\alpha \leq U'$  for all short  $\alpha \in \Sigma^+ \setminus \Delta$  different from  $\alpha_2 + \alpha_3$ .

*Case 4.  $\Sigma \cong G_2$ .*

We let  $\Delta = \{\gamma, \delta\}$  with  $\gamma$  long. From the second formula in (c) we get  $U_{2\gamma+3\delta} \leq U'$ . If  $p > 3$  (resp.  $p = 2$  resp.  $p = 3$ ), then the third formula in (c) gives the following congruences mod  $U_{2\gamma+3\delta}$  (which is a central subgroup of  $U$ ):  $[U_{\gamma+\delta}, U_\delta] \equiv U_{\gamma+2\delta} U_{\gamma+3\delta}$  (resp.  $\equiv U_{\gamma+3\delta}$  resp.  $\equiv U_{\gamma+2\delta}$ ). So if  $p > 3$  then  $U_{\gamma+2\delta}, U_{\gamma+3\delta}, U_{2\gamma+3\delta} \leq U'$ , and then also  $U_{\gamma+\delta} \leq U'$  by the first formula in (c). This settles the case  $p > 3$ . If  $p = 2$  (resp.  $p = 3$ ), then  $U_{\gamma+3\delta}, U_{2\gamma+3\delta} \leq U'$  (resp.  $U_{\gamma+2\delta}, U_{2\gamma+3\delta} \leq U'$ ), and if additionally  $q = 2^n > 2$  (resp.  $q = 3^n > 3$ ) then the first formula in (c) gives also  $U_\alpha \leq U'$  for the remaining two  $\alpha \in \Sigma^+ \setminus \Delta$ . ■

For any  $J \subset \Pi$ , set  $D_J = (G_J U_J)'$ . This group will play a key role in the proof of Proposition 1 below.

**LEMMA 2.** *Let  $J = \Pi \setminus \{i\}$  for some  $i \in \Pi$ . Assume rank  $G > 2$  if  $q \leq 3$ . Then  $U_\alpha \leq D_J$  for all  $\alpha \in \Sigma^+ \setminus \Delta$ , unless  $q = 2$ ,  $\Sigma \cong B_l$  or  $C_l$ , and  $i = l - 2$ .*

*Proof.* By Lemma 1 (and by using the isomorphism  $B_l(2) \rightarrow C_l(2)$ ) it only remains to consider the following two cases:

*Case A.  $q = 2$ ,  $\Sigma \cong B_l$  ( $l \geq 3$ ),  $i \neq l - 2$ ,  $\alpha = \alpha_{l-1} + \alpha_l$  or  $\alpha_{l-1} + 2\alpha_l$ .*

Since  $i \neq l - 2$ , there is  $n \in G_J$  representing the Weyl reflection  $r := r_{\alpha_{l-2}}$ . Then  $r(\alpha)$  is different from the two exceptional roots in Lemma 1(i), and lies in  $\Sigma^+ \setminus \Delta$ . Hence  $U_{r(\alpha)} \leq U'$  by Lemma 1, and so

$$U_\alpha = n^{-1} U_{r(\alpha)} n \leq n^{-1} U' n \leq D_J.$$

*Case B.  $q = 2$ ,  $\Sigma \cong F_4$ ,  $\alpha = \alpha_2 + \alpha_3$  or  $\alpha_2 + 2\alpha_3$ .*

In this case there is  $n \in G_J$  representing either  $r_{\alpha_1}$  or  $r_{\alpha_4}$ . Now argue as in Case A. ■

LEMMA 3. Let  $J = \Pi \setminus \{i\}$  for some  $i \in \Pi$ . If  $q \leq 3$  (resp.  $q = 2$ ) assume that the diagram of  $\Delta_J$  has no isolated node (resp. no connected component of type  $B_2$ ). Then  $U \leq D_J$ .

*Proof.* The hypothesis guarantees that  $G_J$  is perfect (see [St, Lemma 32']). Thus  $U_{\alpha_j} \leq G_J \leq D_J$  for all  $j \in J$ . Furthermore  $U_\alpha \leq D_J$  for all  $\alpha \in \Sigma^+ \setminus \Delta$  by Lemma 2. Thus it suffices to show  $U_{\alpha_i} \leq D_J$ . Choose  $j \in J$  with  $(\alpha_j, \alpha_i) \neq 0$ , and choose  $n \in G_J$  representing  $r := r_{\alpha_j}$ . Then  $r(\alpha_i) \in \Sigma^+ \setminus \Delta$ , hence  $U_{\alpha_i} = n^{-1}U_{r(\alpha_i)}n \leq n^{-1}D_Jn = D_J$ . ■

Now let  $K$  be an algebraic closure of  $k$ . (In fact, our results remain true with  $K$  being any overfield of  $k$ , since  $k$  is a splitting field for  $G$ ). By [St, Th. 43] the equivalence classes of irreducible  $KG$ -modules  $V$  are classified by dominant weights  $\lambda = \sum_{i=1}^l m_i \lambda_i$  with  $0 \leq m_i \leq q-1$ . The  $\lambda$  corresponding to  $V$  will be called the highest weight of  $V$ . This correspondence is "inductive" in the following sense: For any  $J \subset \Pi$  let  $V_J$  denote the fixed space in  $V$  of  $U_J$ . Then  $V_J$  is an irreducible module for  $G_J$  (by [S]), and if  $J$  is non-empty then the highest weight of  $V_J$  (as  $KG_J$ -module) is  $\sum_{j \in J} m_j \lambda'_j$  (where the  $\lambda'_j$  ( $j \in J$ ) are the fundamental weights of  $\Sigma_J$  corresponding to  $\Delta_J$ ). If  $J = \emptyset$ , i.e.,  $P_J = B$ , then  $V_B := V_\emptyset = C_V(U)$  is 1-dimensional and  $T$  acts on it via  $\lambda$ .

From now on  $V$  denotes an irreducible  $KG$ -module with highest weight  $\lambda$ . We consider a non-splitting extension

$$0 \longrightarrow K \longrightarrow E \xrightarrow{\pi} V \longrightarrow 0$$

as in the Introduction. For each  $J \subset \Pi$  let  $E_J$  denote the fixed space in  $E$  of  $U_J$ , and set  $E_B := E_\emptyset = C_E(U)$ . As explained in the Introduction, we want to study the "highest weight module condition" (\*) for  $E$ . Note that (\*) is equivalent to

$$\dim E_B > 1. \quad (*)'$$

We are now ready to give our first criterion for (\*). Recall that  $l = \text{rank } G > 1$ .

PROPOSITION 1. Suppose  $\lambda$  is a multiple of a fundamental weight  $\lambda_i$ , and set  $J := \Pi \setminus \{i\}$ . If  $q \leq 3$  (resp.  $q = 2$ ) assume that the diagram of  $\Delta_J$  has no isolated node (resp. no connected component of type  $B_2$ ). Then (\*) holds for each non-splitting extension  $E$  of  $V$  over  $K$ .

*Proof.* The parabolic  $P_J$  fixes the 1-space  $V_B (= V_J)$  in  $V$ . Thus  $P_J$  also fixes the 2-space  $Y := \pi^{-1}(V_B)$  in  $E$ , and acts triangularly in  $Y$ . Since the

subgroup  $G_J U_J$  of  $P_J$  is generated by  $p$ -elements, it acts in  $Y$  as a subgroup of

$$\left\{ \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} : x \in k \right\}.$$

So  $D_J = (G_J U_J)'$  acts trivially in  $Y$ . But  $U \leq D_J$  by Lemma 3, hence  $Y \subset C_E(U) = E_B$ . This proves  $(*)'$ , hence  $(*)$ . ■

*Remark.* The additional hypothesis for small  $q$  cannot be dropped completely (see Remark 2 below). For fundamental  $\lambda$ , we will settle the cases excluded in Proposition 1. (Note that in these exceptions we have  $q \leq 3$ , hence  $\lambda = \lambda_i$  or  $\lambda = 2\lambda_i$ , the latter only occurring for  $q = 3$ ).

LEMMA 4. *If  $\dim E_J > 1$  for some  $J \subset \Pi$ , we have the exact sequence*

$$0 \rightarrow K \rightarrow E_J \rightarrow V_J \rightarrow 0$$

*of  $L_J$ -modules.*

*Proof.* The hypothesis implies that  $\pi(E_J)$  is a non-zero  $L_J$ -submodule of  $V_J$ , hence equals  $V_J$  (since  $V_J$  is irreducible under  $L_J$ ). ■

PROPOSITION 2. *Suppose  $\lambda$  is a fundamental weight  $\lambda_i$ , and exclude the cases*

- (a)  $q = 2$ ,  $\Sigma \cong A_2$ .
- (b)  $q = 2$ ,  $\Sigma \cong A_3$ , and  $i = 2$ .
- (c)  $q = 3$ ,  $\Sigma \cong B_2$ , and  $i = 1$ .

*Then  $(*)$  holds for each non-splitting extension  $E$  of  $V$  over  $K$ .*

*Proof.* We assume the hypothesis of the Proposition. Additionally, we may assume that we are in one of the cases excluded in Proposition 1:

(1) Either  $q \leq 3$  and the diagram of  $A_{\Pi \setminus \{i\}}$  has an isolated node, or  $q = 2$  and the diagram has a connected component of type  $B_2$ .

The strategy of proof is to reduce to small rank cases, where we can apply the results of [JP]:

(2) Condition  $(*)$  holds if  $\lambda$  is "minimal" in the sense of [JP].

Namely, in this case  $H^1(G, V)$  is known: First, exclude the case  $q = 2$ ,  $\Sigma \cong C_l$  ( $l \geq 3$ ). Since we also excluded the cases (a), (b), (c), it follows from [JP, Sect. 3, 5, and 6C] that if  $\lambda$  is minimal then  $E$  is a quotient of the

Weyl module  $W_\lambda$  corresponding to  $V$ . But  $W_\lambda$  is generated by a  $B$ -semi-invariant, hence so is  $E$ .

Now assume  $q=2$ ,  $\Sigma \cong C_l$  ( $l \geq 3$ ). Then only  $\lambda_1$  and  $\lambda_2$  are minimal. If  $\lambda = \lambda_2$  then again  $E$  is a quotient of  $W_\lambda$  (which follows by comparing [JP, 6B] and [CPS, table on p. 173]). If  $\lambda = \lambda_1$  then we reduce to the case  $\Sigma \cong B_l$ ,  $\lambda = \lambda_1$  by the exceptional isomorphism  $B_l(2) \rightarrow C_l(2)$  (cf. [CPS, p. 182]) and use that  $\lambda_1$  is also minimal for type  $B_l$  (but not  $\lambda_2$ ). This completes the proof of (2).

In the rest of the proof, we check all irreducible root systems individually. For the list of minimal weights, see [JP, 6B].

*Case 1.*  $\Sigma \cong A_l$  ( $l < 5$ ).

In this case all fundamental weights are minimal, hence (2) applies. (In fact, the restriction  $l < 5$  is unnecessary, but I want to stress that we need the results of [JP] only in small rank cases.)

*Case 2.*  $\Sigma \cong G_2$ .

In this case  $\lambda_2$  is minimal. The case  $\lambda = \lambda_1$  reduces to the case  $\lambda = \lambda_2$  if  $q=3$  (by the exceptional graph automorphism of  $G_2(3)$ ). Finally, the case  $\lambda = \lambda_1$ ,  $q=2$  cannot occur here, since in this case  $H^1(G, V^*) = H^1(G, V) = 0$  (see the Appendix), contradicting the existence of non-splitting  $E$ .

*Case 3.*  $\Sigma \cong B_2$ .

Both fundamental weights are minimal, so again (2) applies.

*Case 4.*  $\Sigma \cong C_3$ .

By (1) we have  $\lambda = \lambda_1$  or  $\lambda_2$ . But both  $\lambda_1$  and  $\lambda_2$  are minimal.

*Case 5.*  $\Sigma \cong B_3$ .

If  $q=3$  then  $\lambda = \lambda_2$  by (1); in this case again  $H^1(G, V^*) = H^1(G, V) = 0$  (see the Appendix). The case  $q=2$  reduces to Case 4, since  $B_3(2) \cong C_3(2)$ .

*Case 6.*  $\Sigma \cong F_4$ .

By (1) we have  $\lambda = \lambda_2$  or  $\lambda_3$ . First assume  $\lambda = \lambda_2$ . Since the group  $G_{\{3,4\}} \cong A_2(q)$  is perfect, it acts trivially in  $Y = \pi^{-1}(V_B)$  (by the same argument as in the proof of Proposition 1). Thus  $U_{\alpha_4} \leq G_{\{3,4\}}$  acts trivially in  $Y$ . By Lemma 2, also  $U_\alpha$  acts trivially in  $Y$  for each  $\alpha \in \Sigma^+ \setminus \Delta$ . Setting  $\Omega := \{1, 2, 3\}$ , it follows that  $U_\Omega$  acts trivially in  $Y$ , hence  $\dim E_\Omega \geq \dim Y > 1$ . By Lemma 4, we get the extension

$$0 \rightarrow K \rightarrow E_\Omega \rightarrow V_\Omega \rightarrow 0 \quad (3)$$



of  $G_\Omega$ -modules. Since  $G_\Omega \cong B_3(q)$ , we can invoke Case 5 to conclude that  $E_\Omega$  satisfies the analogue of  $(*)'$ :

$$\dim C_{E_\Omega}(U \cap G_\Omega) > 1.$$

(Note that this trivially holds if  $E_\Omega$  splits over  $K$ .)

Then clearly this centralizer equals  $Y$ , so now  $Y$  is centralized by  $U \cap G_\Omega$ . Above we saw that  $Y$  is centralized by  $U_\Omega$ . It follows that  $Y$  is centralized by  $U = (U \cap G_\Omega)U_\Omega$ . Thus  $Y \leq E_B$ . This proves  $(*)'$ , hence  $(*)$ , if  $\lambda = \lambda_2$ .

The case  $\lambda = \lambda_3$  is analogous (invoking Case 4 instead of Case 5).

*Case 7.*  $\Sigma \cong A_l$  ( $l \geq 5$ ),  $D_l$  ( $l \geq 4$ ) or  $E_l$  ( $l = 6, 7, 8$ ).

In the case  $D_4$  we have  $\lambda = \lambda_2$  by (1), so  $\lambda$  is minimal. Now assume  $\Sigma \neq D_4$ . Then by (1), the diagram of  $\Delta_{\Pi \setminus \{i\}}$  has exactly one or two isolated nodes; let  $I' \subset \Pi$  be the set of  $j$  such that  $\alpha_j$  corresponds to such an isolated node. Set  $I := I' \cup \{i\}$ . Then  $G_{\Pi \setminus I}$  is perfect, and it follows as in Case 6 that  $U_I$  acts trivially in  $Y$ .

Now choose  $\Omega$  with  $I \subset \Omega \subset \Pi$  such that  $\Sigma_\Omega \cong A_4$  or  $D_4$ . Then  $\dim E_\Omega \geq \dim E_I > 1$ , so again we get the extension (3) of  $G_\Omega$ -modules. Since  $G_\Omega \cong A_4(q)$  or  $D_4(q)$ , we can invoke Case 1 (resp. the above case of  $D_4$ ) to conclude that  $E_\Omega$  satisfies the analogue of  $(*)'$ . From this it follows as in Case 6 that  $E$  satisfies  $(*)'$ .

*Case 8.*  $\Sigma \cong B_l$  or  $C_l$  ( $l \geq 4$ ).

By (1) we have  $\lambda = \lambda_2$ ,  $\lambda_{l-2}$  or  $\lambda_{l-1}$ . The weight  $\lambda_2$  is minimal for type  $C_4$ , and if  $q = 2$  then the case  $\Sigma \cong B_4$ ,  $\lambda = \lambda_2$  reduces to that as above. So we can exclude the case  $l = 4$ ,  $\lambda = \lambda_2$ ,  $q = 2$ .

Now set  $I := \{1, 2\}$  if  $\lambda = \lambda_2$ ,  $I := \{l-1, l\}$  if  $\lambda = \lambda_{l-1}$ , and  $I := \{l-2, l-1, l\}$  if  $\lambda = \lambda_{l-2}$  and  $l > 4$ . Then  $G_{\Pi \setminus I}$  is perfect (by the exclusion in the previous paragraph). Thus  $G_{\Pi \setminus I}$  acts trivially in  $Y$ , and so the same is true for each  $U_\alpha$  with  $\alpha \in \Sigma_{\Pi \setminus I}$ . If  $I = \{1, 2\}$  it follows by Lemma 1 that also for each  $\alpha \in \Sigma^+ \setminus \Delta$ , the group  $U_\alpha$  acts trivially in  $Y$ ; then also  $U_I$  acts trivially in  $Y$ . In the other two cases, the exceptional roots from Lemma 1(i), (i)' lie in  $\Sigma_I$ , hence Lemma 1 implies that for all  $\alpha \in \Sigma^+ \setminus (\Delta \cup \Sigma_I)$ , the group  $U_\alpha$  acts trivially in  $Y$ . Thus in all cases,  $U_I$  acts trivially in  $Y$ .

Now we choose  $\Omega$  with  $I \subset \Omega \subset \Pi$  such that  $\Sigma_\Omega \cong A_2$  if  $l = 4$ ,  $\lambda = \lambda_2$ ,  $q = 3$ ,  $\Sigma_\Omega \cong A_4$  if  $l > 4$ ,  $\lambda = \lambda_2$ , and  $\Sigma_\Omega \cong B_3$  or  $C_3$  otherwise. By the previous paragraph,  $\dim E_\Omega \geq \dim E_I > 1$ . Thus we get again the extension (3), which satisfies the analogue of  $(*)'$  by Case 1, 4 and 5. Now it follows as in Case 6 (and 7) that  $E$  satisfies  $(*)'$ , hence  $(*)$ . ■

*Remark 1.* In the cases (a), (b), (c), we have  $\dim H^1(G, V^*) = 1$  by [JP], and *none* of the non-splitting extensions of  $V$  over  $K$  satisfies (\*) (see Remark 2 below).

Condition (\*) implies that  $E$  is a quotient of the induced module  $I(V) := \text{Ind}_B^G(V_B)$ . This is not yet very helpful for the actual computation of  $H^1(G, V^*)$ , since the structure of  $I(V)$  is not transparent. But in many cases, we can replace  $I(V)$  by the much smaller module  $\tilde{V}$ , defined as follows: Let  $f: I(V) \rightarrow V$  be the canonical  $KG$ -module map (extending the inclusion map  $V_B \rightarrow V$ ). For each  $j \in \Pi$ , let  $M_j$  be the  $G_{\{j\}}$ -submodule of  $I(V)$  generated by  $V_B$ . Now let  $M$  be the  $G$ -submodule of  $I(V)$  generated by all the  $M_j \cap \ker(f)$  ( $j \in \Pi$ ), and define  $\tilde{V}$  to be  $I(V)/M$ . (Then  $\tilde{V} = H_0(\mathcal{U}_i)$  in the set-up of [RS-2, 3B]; for the definition adopted here, see [R], where it is also proved that  $\tilde{V}$  has a unique maximal submodule). More geometrically,  $\tilde{V}$  can be viewed as the  $K$ -vector space abstractly generated by the  $G$ -conjugates of the 1-space  $V_B$ , subject to those relations that hold inside the  $G$ -conjugates of the spaces  $V_{\{j\}}$ . In fact,  $\tilde{V}$  has explicitly been computed in several cases by geometric methods (see below).

LEMMA 5. Assume (\*) holds, and the extension

$$0 \rightarrow K \rightarrow E_{\{j\}} \rightarrow V_{\{j\}} \rightarrow 0$$

of  $G_{\{j\}}$ -modules splits for every  $j \in \Pi$  with  $\dim V_{\{j\}} > 1$ . Then  $E$  is a quotient of  $\tilde{V}$ .

*Proof.* First note that the above extension exists by Lemma 4, and splits automatically if  $\dim V_{\{j\}} = 1$ . (Namely in this case,  $E_{\{j\}} = E_B$ , and so  $U \cap G_{\{j\}}$ , hence  $G_{\{j\}}$ , acts trivially in  $E_{\{j\}}$ ). Clearly, the extension splits also under  $L_{\{j\}}$ .

By (\*) there is a  $KB$ -module map  $\iota: V_B \rightarrow E$  such that  $\pi\iota = \text{id}_{V_B}$ . (Recall that  $\pi$  is the map  $E \rightarrow V$ ). We can choose  $\iota$  such that

(1) For each  $j \in \Pi$ , the  $G_{\{j\}}$ -submodule  $\bar{V}_{\{j\}}$  of  $E_{\{j\}}$  generated by  $\iota(V_B)$  splits  $E_{\{j\}}$  over  $\ker(\pi)$ :

Namely, if  $\lambda(T) \neq 1$  then  $\iota(V_B)$  is the unique 1-dimensional  $KB$ -submodule of  $E_{\{j\}}$  different from  $\ker(\pi)$ , hence (1) follows from the fact that  $E_{\{j\}}$  splits under  $L_{\{j\}}$ . Now assume  $\lambda(T) = 1$ . Then  $\lambda = \sum_{j=1}^l m_j \lambda_j$  with  $m_j = 0$  or  $q-1$  (since  $1 = \lambda(h_{\alpha_j}(s)) = s^{\langle \lambda, \alpha_j \rangle} = s^{m_j}$  for each  $s \in k^*$ ). Let  $J$  be the set of those  $j$  with  $m_j = q-1$ . Then  $J$  is non-empty (since otherwise  $\lambda = 0$  and (\*) forces  $B$ , hence  $G$ , to act trivially in  $E$ , which contradicts our standing hypothesis that  $E$  is non-splitting). Now  $V_J$  is the Steinberg module of  $G_J$ , hence is projective. So  $E_J$  splits over  $\ker(\pi)$  as  $G_J$ -module. Thus we can choose  $\iota$  so

that  $\iota(V_B)$  lies in the unique  $G_J$ -submodule of  $E_J$  isomorphic to  $V_J$ . Then (1) is clear for  $j \in J$ . But if  $j \notin J$  then  $G_{\{j\}}$  acts trivially in  $E_{\{j\}}$ , and so (1) holds again.

Let  $h: I(V) \rightarrow E$  be the  $KG$ -module map extending  $\iota$ . Then  $\pi h = f$ . Thus  $h$  maps  $\ker(f)$  into  $\ker(\pi)$ . On the other hand,  $h$  maps  $M_j$  into  $\tilde{V}_{\{j\}}$  (by definition of these modules). By (1), it follows that  $h$  vanishes on  $M_j \cap \ker(f)$  for each  $j \in \Pi$ . Thus the map  $h: I(V) \rightarrow E$  vanishes on  $M$ , hence factors through  $\tilde{V}$ . ■

Once (\*) is known to hold, the hypothesis of Lemma 5 can easily be verified in many cases, since the extension theory for irreducible modules of  $G_{\{j\}} \cong SL_2(q)$  is known [AJL]. For the case of fundamental modules, we need only the elementary fact that the 1-cohomology of  $SL_2(q)$  in its natural module vanishes unless  $q = 2^n > 2$ :

**COROLLARY 1.** *Suppose  $q = p$  is a prime, or  $\lambda$  is a fundamental weight. Then each non-splitting extension  $E$  of  $V$  over  $K$  satisfying (\*) is a quotient of  $\tilde{V}$ .*

*Proof.* If (\*) holds for  $E$ , then the analogue of (\*)' holds for each  $E_{\{j\}}$ ; hence if  $E_{\{j\}}$  is non-splitting then it is a quotient of  $\text{Ind}_{G_{\{j\}} \cap B}^{G_{\{j\}}}(V_B)$ . But it is a well known fact that a  $K[SL_2(p)]$ -module induced from a 1-dimensional module  $Z$  of a Borel subgroup has exactly two composition factors, neither of which is trivial unless  $Z$  is trivial; and if  $Z$  is trivial then the induced module is semisimple. Thus if  $q = p$  is a prime, the extension in Lemma 5 splits for each  $j \in \Pi$ , proving the claim in this case.

Now suppose  $\lambda = \lambda_i$  is fundamental. Then  $V_{\{j\}} = V_B$  for each  $j \neq i$ , and  $V_{\{i\}}$  is 2-dimensional, affording the natural module of  $G_{\{i\}} \cong SL_2(q)$ . Thus  $H^1(G_{\{i\}}, V_{\{i\}}^*) \cong H^1(G_{\{i\}}, V_{\{i\}}) = 0$  unless  $q = 2^n > 2$ . By Lemma 5, it only remains to show that the extension

$$(1) \quad 0 \rightarrow K \rightarrow E_{\{i\}} \rightarrow V_{\{i\}} \rightarrow 0$$

of  $G_{\{i\}}$ -modules splits for  $q = 2^n > 2$ .

Exclude for a moment the case that  $q = 4$ ,  $\Sigma \cong G_2$  and  $\alpha_i$  is long. Then we have:

$$(2) \quad \text{There is } t \in T \text{ with } \lambda_i(t) \neq 1, \alpha_i(t) = 1.$$

This implies the claim as follows: Since  $T$  acts in  $V_{\{i\}}$  with weights  $\lambda_i, \lambda_i - \alpha_i$  (restricted to  $T$ ), it follows that  $t$  acts in  $V_{\{i\}}$  as scalar multiplication by  $\lambda_i(t) \neq 1$ . Furthermore  $t$  centralizes  $G_{\{i\}}$  (since  $\alpha_i(t) = 1$ ). Hence the  $\lambda_i(t)$ -eigenspace of  $t$  in  $V_{\{i\}}$  splits (1).

Now we prove (2) (under the hypothesis that  $q = 2^n > 2$ , and  $q > 4$  if

$\Sigma \cong G_2$  and  $\alpha_i$  is long). Choose  $j \in I \setminus \{i\}$  with  $(\alpha_i, \alpha_j) \neq 0$ , and take  $t$  of the form  $t = h_{\alpha_i}(s_i)h_{\alpha_j}(s_j)$  with  $s_i, s_j \in k^*$ . Then

$$\begin{aligned}\lambda_i(t) &= s_i^{\langle \lambda_i, \alpha_i \rangle} s_j^{\langle \lambda_i, \alpha_j \rangle} = s_i \\ \alpha_i(t) &= s_i^{\langle \alpha_i, \alpha_i \rangle} s_j^{\langle \alpha_i, \alpha_j \rangle} = s_i^2 s_j^{-\varepsilon},\end{aligned}$$

where  $\varepsilon = -\langle \alpha_i, \alpha_j \rangle = 1, 2$ , or  $3$ . If  $\varepsilon = 1$  or  $2$ , then there is  $s_j \in k^*$  with  $s_j^\varepsilon \neq 1$  (since  $q = 2^n > 2$ ). The case  $\varepsilon = 3$  occurs only if  $\Sigma \cong G_2$  and  $\alpha_i$  is long; in this case we have  $q > 4$ , and so there is  $s_j \in k^*$  with  $s_j^\varepsilon = s_j^3 \neq 1$ . Then in any case, we have  $s_i \in k^*$  with  $s_i^2 = s_j^\varepsilon \neq 1$ , hence also  $s_i \neq 1$  (since  $p = 2$ ). Now  $t = h_{\alpha_i}(s_i)h_{\alpha_j}(s_j)$  satisfies (2).

It remains to settle the case that  $q = 4$ ,  $\Sigma \cong G_2$  and  $\alpha_i$  is long. Then  $V$  is the 14-dimensional adjoint module of  $G_2(4)$ ; under  $G_0 := \langle U_\alpha : \alpha \in \Sigma \text{ long} \rangle \cong A_2(4)$  it decomposes as  $V = A \oplus V_3 \oplus \bar{V}_3$ , where  $A$  is the (irreducible) 8-dimensional adjoint module of  $A_2(4)$ , and  $V_3, \bar{V}_3$  are the natural 3-dimensional module and its dual (see the Appendix). Thus  $H^1(G_0, V^*) \cong H^1(G_0, V) \cong H^1(G_0, A) \oplus H^1(G_0, V_3) \oplus H^1(G_0, \bar{V}_3) = 0$  (by [CPS]), hence  $E$  splits as  $KG_0$ -module. But  $G_{\{i\}} \leq G_0$ ; so  $E$  splits under  $G_{\{i\}}$ , hence the same is true for  $E_{\{i\}}$ . ■

**THEOREM 1.** *Suppose that  $\text{rank } G > 1$ , and the irreducible  $KG$ -module  $V$  is fundamental with highest weight  $\lambda_i$ , excluding cases (a)–(c) of Proposition 2. Then each non-splitting extension  $E$  of  $V$  over  $K$  is a quotient of  $\tilde{V}$ .*

*Proof.* Proposition 2 and Corollary 1.

**Remark 2.** In cases (a)–(c) we have  $\tilde{V} \cong V$  by [RS-1], and  $\dim H^1(G, V^*) = 1$  by [JP, 6C]. So these cases actually give rise to counterexamples.

**Remark 3.**  $\tilde{V}$  has been calculated essentially for the modules  $V$  minimal in the sense of [RS-1] (in that paper) and for the adjoint modules not of type  $C_l$  (in [SV], [V-1]). In all these cases except for the adjoint module of type  $A_l$ , the module  $V$  is fundamental and so Theorem 1 yields  $H^1(G, V^*)$  (with the exceptions from (a)–(c)). (We could easily adapt our method to include the adjoint module of  $A_l$ , but this does not seem worthwhile, since the 1-cohomology of this module is known by [JP]).

Most of these cohomology groups have already been calculated in [CPS] and [JP], except for the adjoint modules of type  $B_l, C_l, G_2$ , and  $F_4$ . An announced result of Hertz [He] (proofs have never appeared) covers also the case of  $F_4$ .

In [V-2] the present author develops another approach to the 1-cohomology of the adjoint modules, which is more general in that it applies also to the case of infinite ground fields (as in [He]) and covers the

case  $C_l$  as well. However, that approach breaks down for small fields ( $q \leq 9$ ). By contrast, the methods of the present paper are primarily aimed at small fields.

Summarizing, we now know the 1-cohomology of all irreducible modules of adjoint type, except type  $C_l$  ( $l > 2$ ) for  $q = 3, 5, 9$ . Thereby, by "irreducible module of adjoint type" we mean the irreducible  $KG$ -module  $V$  whose highest weight is the highest root of  $\Sigma$  (i.e.,  $V$  is the unique irreducible quotient of the usual adjoint module).

**COROLLARY 2.** *Assume  $\Sigma$  is not of type  $A_l$ ,  $C_l$ , or  $B_2$ . Then for the irreducible  $KG$ -module  $V$  of adjoint type, we have  $H^1(G, V) = 0$  unless  $p = 2$ ,  $\Sigma \cong B_l$  ( $l > 2$  even) or  $p = 3$ ,  $\Sigma \cong E_6$  or  $p = 2$ ,  $\Sigma \cong E_7$  or  $p = 2$ ,  $\Sigma \cong D_l$  ( $l > 3$ ). In these exceptional cases we have  $\dim H^1(G, V) = 1$  except in the last one for  $l$  even, where  $\dim H^1(G, V) = 2$ .*

*Proof.* Let  $L$  denote the usual adjoint module of  $G$ , i.e., the Lie algebra of the (simply-connected) algebraic group  $G_K$  enclosing  $G$  (whereby  $G$  acts on  $L$  in the restriction of the adjoint representation of  $G_K$ ). Then  $V$  is the unique irreducible quotient of  $L$  (and  $L$  is the Weyl-module corresponding to  $V$ ); by [V-1, Cor. 2] the module  $\tilde{V}$  is isomorphic to  $L$  (since  $\Sigma$  is not of type  $C_l$  or  $B_2$ ). (Strictly speaking, in [V-1] the analogue of  $\tilde{V}$  over  $k$  (instead of  $K$ ) is studied, but the transition is immediate).

The highest weight of  $V$  is the highest root of  $\Sigma$ , hence is fundamental in the cases under consideration (see [Bou, appendix]). Also  $V \cong V^*$ , hence by Theorem 1 the dimension of  $H^1(G, V)$  can be read off from  $\tilde{V} \cong L$ : It equals the multiplicity of the trivial representation in  $M/\text{rad}(M)$ , where  $M$  is the unique maximal submodule of  $L$ . This multiplicity is given by [H] (and [Hur]), where the full submodule structure of  $L$  is determined. ■

### 3. MORE ON THE CONDITION (\*)

In this section,  $l = \text{rank } G \geq 1$  (so we now include the case  $l = 1$ ). Otherwise, we keep all notations and conventions from Sections 1 and 2. In particular,  $V$  is an irreducible  $KG$ -module with highest weight  $\lambda$ , and

$$0 \longrightarrow K \longrightarrow E \xrightarrow{\pi} V \longrightarrow 0$$

is a non-splitting extension of  $KG$ -modules. We set  $Y := \pi^{-1}(V_B)$  (a 2-dimensional subspace of  $E$ ).

LEMMA 6. If  $U_\alpha$  acts non-trivially in  $Y$  for some  $\alpha \in \Sigma$ , then

$$\lambda + p^r \alpha \in (q-1)X$$

for some integer  $r$  with  $0 \leq r \leq n-1$ .

*Proof.* Fix  $\alpha \in \Sigma$ , and pick a basis  $y, z$  of  $Y$  consisting of weight vectors for  $T$ , with  $z \in \ker(\pi)$ . Then  $y$  is a vector of weight  $\lambda$ . Furthermore, there is an endomorphism  $f$  of the additive group of  $k$  with  $x_\alpha(a) \cdot y = y + f(a)z$  for each  $a \in k$ . Since we assume that  $U_\alpha$  acts non-trivially in  $Y$ , we have  $f \neq 0$ .

For all  $t \in T$ ,  $a \in k$  we get  $y + f(\alpha(t)a)z = x_\alpha(\alpha(t)a) \cdot y = [tx_\alpha(a)t^{-1}] \cdot y = [tx_\alpha(a)] \cdot \lambda(t^{-1})y = y + \lambda(t)^{-1}f(a)z$ , thus

$$f(\alpha(t)a) = \lambda(t)^{-1}f(a) \quad \text{for all } t \in T, \quad a \in k. \quad (1)$$

If  $f(1) = 0$  then  $f(s^2) = f(\alpha(h_\alpha(s))) = \lambda(h_\alpha(s))^{-1}f(1) = 0$  for all  $s \in k^*$ ; thus  $f = 0$ , since  $f(a) = f((a/2 + 1)^2) - f(a^2/4) - f(1^2) = 0$  if  $p \neq 2$ , and  $k = k^2$  if  $p = 2$ . But we have  $f \neq 0$ , hence  $f(1) \neq 0$ . This allows us to assume  $f(1) = 1$  (replacing  $z$  by  $f(1)z$ ). Then  $f(\alpha(t)) = \lambda(t)^{-1}f(1) = \lambda(t)^{-1}$ , hence  $f(\alpha(t)a) = \lambda(t)^{-1}f(a) = f(\alpha(t))f(a)$  for all  $t \in T$ ,  $a \in k$ . Thus

$$f(ab) = f(a)f(b) \quad (2)$$

if either  $a$  or  $b$  is a square. In fact, (2) holds for all  $a, b \in k$ : This is clear if  $p = 2$ ; if  $p \neq 2$ , then  $f(ab) = f([(a/2 + 1)^2 - a^2/4 - 1]b) = f((a/2 + 1)^2)f(b) - f(a^2/4)f(b) - f(1)f(b) = f(a)f(b)$ .

Thus  $f$  is a ring homomorphism, and since  $f \neq 0$  it follows that  $f$  is an automorphism of  $k$ . Since  $k$  is the field with  $q = p^n$  elements, there is  $r$  with  $0 \leq r \leq n-1$  such that  $f(a) = a^{p^r}$  for all  $a \in k$ . Now (1) gives  $\lambda(t)^{-1} = f(\alpha(t)) = \alpha(t)^{p^r}$ . Thus  $\lambda + p^r \alpha$  vanishes on  $T$ , hence lies in  $(q-1)X$ . ■

The extension theory for irreducible modules of  $SL_2(q)$  is known by [AJL]. In the case  $G = SL_2(q)$ , this allows us to determine all extensions  $E$  satisfying (\*). This will be the base for the higher rank cases.

LEMMA 7. Suppose  $G = SL_2(q)$  (i.e.,  $l = 1$ ).

- (1) If  $q = p$  then (\*) never holds.
- (2) If  $q \neq p$  then (\*) always holds.

Furthermore, in the case  $q = p > 2$  the non-splitting extension  $E$  can only exist if  $\lambda = (p-3)\lambda_1$ .

*Proof.* (1) See the case  $q = p$  of the proof of Corollary 1.

(2) We have  $\lambda = m\lambda_1$  with  $0 \leq m \leq q-1$ . Since we are now in the case  $q = p^n \neq p$ , it follows from [AJL, Cor. 4.5] that the non-splitting extension  $E$  can only exist for the following values of  $m$ :

$$1 + (p-2)p^{n-1}, \quad p^s((p-2)+p) \quad \text{for } 0 \leq s \leq n-2.$$

We first consider the latter case. In this case  $V$  is the  $s$ th-Frobenius twist of the irreducible module with highest weight  $((p-2)+p)\lambda_1$ , hence we may assume  $s=0$ . Then by Steinberg's tensor product theorem,  $V \cong V_{p-2} \otimes V_1^{(p)}$ , where  $V_{p-2}$  is the irreducible module with highest weight  $(p-2)\lambda_1$  and  $V_1^{(p)}$  is the Frobenius twist of the 2-dimensional natural module. Thus  $\dim V = 2(p-1)$ . On the other hand, the Weyl module  $W_\lambda$  corresponding to  $V$  has dimension  $m+1 = 2p-1$ . Thus  $W_\lambda$  is a non-splitting extension of  $V$  over  $K$  (cf. [W, Th. 2D]). Since  $\dim \text{Ext}_{KG}^1(V, K) = 1$  by [AJL, Cor. 4.5], each such non-splitting extension  $E$  is isomorphic to  $W_\lambda$ . But  $W_\lambda$  satisfies (\*), so we are done.

To deal with the remaining case  $m = 1 + (p-2)p^{n-1}$ , we assume by way of contradiction that (\*) does not hold. Then  $U = U_{\alpha_1}$  acts non-trivially in  $Y$ , hence by Lemma 6 we have  $\lambda + p^r\alpha_1 \in (q-1)X$  for some  $r$  with  $0 \leq r \leq n-1$ . Since  $\alpha_1 = 2\lambda_1$ , we get  $m + 2p^r \mid (q-1)$ . Then from  $m + 2p^r = p^n - 2p^{n-1} + 2p^r + 1 \leq p^n + 1$  it follows that  $m + 2p^r = q-1$ , hence  $2p^{n-1} - 2p^r = 2$ . So  $p^{n-1} - p^r = 1$ , which is only possible if  $p=2$ ,  $n=2$ ,  $r=0$ . In this case,  $\dim H^1(G, V) = \dim H^1(SL_2(4), K^2) = 1$ , hence the Frobenius twist of  $E$  is isomorphic to the module of traceless  $2 \times 2$ -matrices, which is a non-splitting extension of  $V_1^{(2)}$  over  $K$  satisfying the analogue of (\*) (cf. [CPSK, 7.6(a)]). This settles the case  $p=2$ ,  $n=2$ ,  $r=0$ , and thereby case (2).

The last assertion in the Lemma follows from the known (elementary) extension theory for irreducible modules of  $SL_2(p)$ , see e.g., [A1, p. 48–49]. (It also follows from Lemma 6). ■

**THEOREM 2.** *Suppose that either  $q$  is not a prime, or  $q > 2$  and  $\langle \lambda, \alpha_j \rangle \neq q-3$  for all  $j \in \Pi$ . Assume  $q > 3$  if  $\Sigma \cong G_2$ . Then (\*) holds for each non-splitting extension  $E$  of  $V$  over  $K$ .*

*Proof.* In view of the extra hypothesis in the case  $G_2$ , Lemma 1 yields  $U_\alpha \leq U'$  for each  $\alpha \in \Sigma^+ \setminus A$ ; hence  $U_\alpha$  acts trivially in  $Y$ . It remains to show that also the  $U_{\alpha_j}$  act trivially in  $Y$ .

So fix  $j \in \Pi$ . We consider the extension

$$0 \rightarrow K \rightarrow \pi^{-1}(V_{\{j\}}) \rightarrow V_{\{j\}} \rightarrow 0 \quad (\text{A})$$

of  $G_{\{j\}}$ -modules. By Lemma 7, this extension always splits in the case  $q = p > 2$ , since then  $V_{\{j\}}$  is the irreducible module for  $G_{\{j\}} \cong SL_2(p)$  of

highest weight  $m_j \lambda'_j$  and  $m_j := \langle \lambda, \alpha_j \rangle \neq p-3$ . But if (A) splits, then clearly  $U_{\alpha_j}$  acts trivially in  $Y$ .

If  $q \neq p$  and the extension (A) does not split, then it satisfies the analogue of (\*) by Lemma 7, which again means that  $U_{\alpha_j}$  acts trivially in  $Y$ . Thus we have shown that  $U_{\alpha_j}$  acts trivially in  $Y$  in any case, which completes the proof of the theorem. ■

*Remark.* The following counterexample shows that the additional hypothesis in the case  $q=p$  is in fact necessary for each prime  $p > 3$  (for  $p \leq 3$ , see Remark 2): If  $q=p > 3$ ,  $\Sigma \cong A_2$  and  $\lambda = (p-3)\lambda_1 + \lambda_2$ , then there exists a non-splitting extension of  $V$  over  $K$  not occurring in the induced module  $\lambda_B^G$  (hence not satisfying (\*)), as remarked in [An, 4.2].

Using Lemma 6 instead of Lemma 7 in the proof of Theorem 2, we get the following

COROLLARY (OF PROOF). *Suppose  $q=p > 2$ , and  $q > 3$  for type  $G_2$ . If*

$$\lambda + \alpha_j \notin (q-1)X$$

*for all  $j \in \Pi$ , then (\*) holds for all non-splitting extensions  $E$  of  $V$  over  $K$ .*

As in Section 1, our criteria for the condition (\*) can be combined with Lemma 5 (and Corollary 1) to yield conditions ensuring that  $E$  is a quotient of  $\tilde{V}$ .

## APPENDIX

Here we provide some computations needed in the proof of Proposition 2. We show  $H^1(G, V) = 0$  if  $G = G_2(2)$  (resp.  $G = B_3(3)$ ) and  $V$  is the irreducible  $KG$ -module of highest weight  $\lambda_1$  (resp.  $\lambda_2$ ). In these cases,  $V$  is the unique irreducible quotient of the adjoint module  $L$  (as in the proof of Corollary 2), and then even  $V = L$  (by [H]).

$L$  comes with a Chevalley basis  $\{E_\alpha, H_\beta : \alpha \in \Sigma, \beta \in \Delta\}$ , such that the action of the generators  $x_\alpha(a)$  of  $G$  on  $L$  is given in terms of this basis by the usual formulas (see, e.g. [H, Sect. 3]). Setting  $L_\alpha := KE_\alpha$ ,  $L_0 := \sum_{\beta \in \Delta} KH_\beta$ , we have in particular:

$$(x_\alpha(a) - 1)L_\beta \subset \sum_{i > 0} L_{i\alpha + \beta} \quad \text{for all } \alpha, \beta \in \Sigma, \quad a \in k. \quad (1)$$

(Thereby of course  $L_{i\alpha + \beta} = 0$  if  $i\alpha + \beta \notin \Sigma \cup 0$ ).

For simplicity we write  $x^\alpha := x_\alpha(1)$  for  $\alpha \in \Sigma$ .

LEMMA A. *Suppose  $G = A_1(q)$  with  $q = 2$  or  $q$  odd, and  $\Sigma = \{\alpha, -\alpha\}$ . If*



$c$  is a cocycle of  $G$  in its adjoint module  $L$  with  $c(U_\alpha) = 0$  and  $c(U_{-\alpha}) \subset L_{-\alpha}$  then  $c = 0$ .

*Proof.* First assume  $q$  is odd. Then  $H^1(A_1(q), L) = 0$ , hence  $c$  is a coboundary, i.e.,  $c(g) = (g - 1)v$  for some  $v \in L$ . Write  $v = tE_\alpha + rE_{-\alpha} + sH_\alpha$  with  $t, r, s \in k$ . Then  $0 = c(x^\alpha) = (x^\alpha - 1)v = rH_\alpha - (2s + r)E_\alpha$ . Hence  $r = s = 0$ . Similarly, the condition  $c(x^{-\alpha}) \subset L_{-\alpha}$  yields  $t = 0$ . So  $c = 0$ .

Now assume  $q = 2$ . Then  $w := x^\alpha x^{-\alpha} x^\alpha$  is an involution, hence  $w \cdot c(w) = c(w)$ . We have  $c(x^{-\alpha}) = aE_{-\alpha}$  for some  $a \in K$ , and so we get  $c(w) = aH_\alpha + aE_\alpha$ . But  $c(w)$  is fixed by  $w$ , which forces  $a = 0$ , hence  $c = 0$ . ■

LEMMA B. Suppose  $q$  is prime, and  $G$  fixes no vector  $\neq 0$  in  $L$ . If  $c$  is a cocycle of  $G$  in  $L$  with  $c(U_\alpha) \subset L_\alpha$  for all  $\alpha \in \Sigma$ , then  $c$  is a coboundary.

*Proof.* First we consider the special case that additionally  $c(U_\alpha) = 0$  for all  $\alpha \in \Delta$ . Then also  $c(U_{-\alpha}) = 0$  (by Lemma A, applied to the subgroup  $\langle U_\alpha, U_{-\alpha} \rangle \cong SL_2(q)$ ), hence  $c = 0$  (since  $G = \langle U_\alpha : \alpha \in \pm \Delta \rangle$ ).

For  $H = \sum_{\beta \in \Delta} a_\beta H_\beta$  ( $a_\beta \in K$ ), we have

$$(x^\alpha - 1)H = \left( - \sum_{\beta \in \Delta} a_\beta c_{\beta\alpha} \right) E_\alpha \quad \text{for all } \alpha \in \Sigma,$$

where the  $c_{\beta\alpha}$  are the Cartan integers (see [H]). If the matrix  $(c_{\beta\alpha})_{\alpha, \beta \in \Delta}$  were singular mod  $p$ , then there would be  $H \neq 0$  as above with  $(x^\alpha - 1)H = 0$  for all  $\alpha \in \Delta \cup -\Delta$ , hence  $H$  would be fixed by  $G$ , contradicting the hypothesis. So the matrix is non-singular mod  $p$ , which means we can choose  $H$  such that  $(x^\alpha - 1)H = c(x^\alpha)$  for all  $\alpha \in \Delta$ . Adjusting  $c$  by the coboundary  $g \rightarrow (g - 1)H$  brings us now back into the above special case. ■

Set  $G_0 := \langle U_\alpha : \alpha \in \Sigma \text{ long} \rangle$ . Then  $G_0 \cong A_2(2)$  (resp.  $G_0 \cong A_3(3)$ ) if  $G = G_2(2)$  (resp.  $G = B_3(3)$ ). Let  $A$  denote the adjoint module of  $G_0$ , naturally embedded into  $L$  (i.e.,  $A = L_0 + \sum_{\alpha \text{ long}} L_\alpha$ ). Using (1) it is easy to see that  $L$  decomposes under  $G_0$  as

$$L = A \oplus V_3 \oplus \bar{V}_3 \quad \text{resp.} \quad L = A \oplus V_6, \quad (2)$$

where  $V_3, \bar{V}_3$  are the natural 3-dimensional module of  $A_2(2)$  and its dual; furthermore  $V_6$  is the 6-dimensional orthogonal module of  $A_3(3) \cong \Omega_6^+(3)$ . (Clearly,  $V_6 = \sum_{\alpha \text{ short}} L_\alpha$ , and  $V_3, \bar{V}_3$  are each spanned by the  $L_\alpha$  where  $\alpha$  runs through an orbit of short roots under the Weyl group of  $G_0$ ).

Now we fix a cocycle  $c$  of  $G$  in  $L$ . If  $G = B_3(3)$  then  $H^1(G_0, L) \cong H^1(G_0, A) \oplus H^1(G_0, V_6) = 0$  by [JP], hence we may assume  $c|_{G_0} = 0$ . For

$G = G_2(2)$ , this will require some more work later. For now, we just assume this condition

$$c|_{G_0} = 0. \quad (3)$$

Pick a short  $\alpha \in \Sigma$ , and let  $C_0$  (resp.  $P_0$ ) denote the centralizer (resp. normalizer) of  $L_\alpha$  in  $G_0$ . Since  $L_\alpha$  is a 1-space in  $V_3$  or  $\bar{V}_3$  (resp. an isotropic 1-space in  $V_6$ ),  $P_0$  is a maximal parabolic subgroup of  $G_0$ ; it follows from (2) that  $P_0$  fixes no other 1-space in  $L$ . Thus  $L_\alpha$  is the full centralizer of  $C_0$  in  $L$  (since this centralizer is invariant under  $P_0$ , and  $P_0/C_0 \cong k^*$ ). Clearly,  $C_0$  centralizes also  $U_\alpha$  (e.g., use the uniqueness in [Hu, Th. 26.3(a)]); by (3) (and (4) below) this implies that  $C_0$  centralizes  $c(U_\alpha)$ , hence  $c(U_\alpha) \subset C_L(C_0) = L_\alpha$ . Thus  $c(U_\alpha) \subset L_\alpha$  for all short roots  $\alpha \in \Sigma$ ; but this trivially holds also for the long roots (by (3)). Hence Lemma B applies, proving that  $c$  is a coboundary.

It remains to justify hypothesis (3) in the case  $G = G_2(2)$ . We proceed similarly as in [J, Ch. 9]. First we record a useful formula from [J, 3.3] (which follows directly from the cocycle condition):

$$\text{For commuting elements } g_1, g_2 \in G \text{ we have } (g_1 - 1)c(g_2) = (g_2 - 1)c(g_1). \quad (4)$$

From now on  $G = G_2(2)$ , and we fix a basis  $\gamma, \delta$  of  $\Sigma$  with  $\gamma$  long. Since  $H^1(G_0, A) = 0$  (by [JP]), we may assume by (2) that  $c(G_0) \subset V_3 \oplus \bar{V}_3$ . So  $c(x^\gamma) = \sum_{\alpha \text{ short}} a_\alpha E_\alpha$  for certain  $a_\alpha \in K$ . Since  $x^\gamma$  is an involution, we have  $x^\gamma \cdot c(x^\gamma) = c(x^\gamma)$ . This implies  $a_\delta = a_{-\gamma-\delta} = 0$  (using (1)).

Let  $c'$  be the coboundary defined by  $c'(g) = (g - 1) \cdot (a_{\gamma+\delta} E_\delta + a_{-\delta} E_{-\gamma-\delta})$ , and set  $c'' := c + c'$ . Then

$$\begin{aligned} c''(x^\gamma) &= c(x^\gamma) + (x^\gamma - 1) \cdot a_{\gamma+\delta} E_\delta + (x^\gamma - 1) \cdot a_{-\delta} E_{-\gamma-\delta} \\ &= \left( \sum_{\alpha \text{ short}} a_\alpha E_\alpha \right) + a_{\gamma+\delta} E_{\gamma+\delta} + a_{-\delta} E_{-\delta} \\ &= a_{\gamma+2\delta} E_{\gamma+2\delta} + a_{-\gamma-2\delta} E_{-\gamma-2\delta}. \end{aligned}$$

Thus, replacing  $c$  by  $c''$ , and setting  $\eta := \gamma + 2\delta$ , we may assume

$$c(x^\gamma) = aE_\eta + bE_{-\eta} \quad (a, b \in K). \quad (5)$$

For later use we note that if for some long  $\gamma' \in \Sigma$  with  $(\gamma, \gamma') < 0$  the original  $c$  satisfied  $c(x^{\gamma'}) = 0$ , then this condition still holds after adjusting with the coboundary  $c'$  (since  $c'(x^{\gamma'}) \subset (x^{\gamma'} - 1) \cdot (L_\delta \oplus L_{-\gamma-\delta}) = 0$  by (1)). By (4) and (5) we get:

$$\begin{aligned} (x^\gamma - 1) \cdot c(x^\eta) &= (x^\eta - 1) \cdot c(x^\gamma) = a(x^\eta - 1)E_\eta \\ &\quad + b(x^\eta - 1)E_{-\eta} = bH_\eta + bE_\eta \end{aligned}$$

If  $b \neq 0$ , it follows by (1) that

$$H_\eta + E_\eta \in (x^\gamma - 1)L \subset L_0 \oplus L_\gamma \oplus L_{\gamma+\delta} \\ \oplus L_{-\gamma-\delta} \oplus L_{2\gamma+3\delta} \oplus L_{-2\gamma-3\delta}.$$

This contradiction proves  $b = 0$ . Similarly, we get  $a = 0$  (replacing  $\eta$  by  $-\eta$  in the above). Now (5) gives:

$$c(x^\gamma) = 0. \quad (6)$$

Since  $\gamma' := \gamma + 3\delta$  is long, we can also adjust  $c$  such that:

$$c(x^{\gamma'}) = 0. \quad (7)$$

As noted above, we can even get (6) and (7) simultaneously. Repeating this argument once more for  $\gamma'' := \gamma - 3\delta$ , we see that we can finally assume  $c$  to vanish on  $x^\gamma$ ,  $x^{\gamma'}$  and  $x^{\gamma''}$ . But these elements generate  $G_0 \cong SL_3(2)$ , and so  $c|_{G_0} = 0$ . This justifies hypothesis (3), and so the proof is complete.

## REFERENCES

- Al. J. L. ALPERIN, "Local Representation Theory," Cambridge Univ. Press, Cambridge, 1986.
- An. H. ANDERSEN, Extensions of simple modules for finite Chevalley groups, *J. Algebra* **111** (1987), 388–403.
- AJL. H. ANDERSEN, J. JORGENSEN AND P. LANDROCK, The projective indecomposable modules of  $SL_2(p^n)$ , *Proc. London Math. Soc.* **46** (1983), 38–52.
- Bou. N. BOURBAKI, "Groupes et algèbres de Lie," Ch. 6, Systèmes de racines," Hermann, Paris, 1968.
- CPS. E. CLINE, B. PARSHALL, AND L. SCOTT, Cohomology of finite groups of Lie type, I, *Publ. Math. IHES* **45** (1975), 169–191.
- CPSK. E. CLINE, B. PARSHALL, L. SCOTT, AND W. VAN DER KALLEN, Rational and generic cohomology, *Invent. Math.* **39** (1977), 143–163.
- He. D. HERTZIG, Cohomology of certain Steinberg groups, *Bull. Amer. Math. Soc.* **75** (1969), 35–36.
- H. H. HISS, Die adjungierten Darstellungen der Chevalley Gruppen, *Arch. Math.* **42** (1984), 408–416.
- Hu. J. E. HUMPHREYS, "Linear Algebraic Groups," 2nd ed., Springer-Verlag, New York, 1981.
- Hur. J. F. HURLEY, Centers of Chevalley algebras, *J. Math. Soc. Japan* **34** (1982), 219–222.
- J. W. JONES, Cohomology of finite groups of Lie type, Thesis, University of Minnesota 1975.
- JP. W. JONES AND B. PARSHALL, On the cohomology of finite groups of Lie type, in "Proc. of the Utah Conf. on Finite Groups," (R. Scott and F. Gross, Eds.), Academic Press, 1976.
- R. M. A. RONAN, Duality for presheaves on chamber systems, and a related chain complex, *J. Algebra* **121** (1989), 263–274.
- RS-1. M. A. RONAN AND S. D. SMITH, Sheaves on buildings and modular representations of Chevalley groups, *J. Algebra* **96** (1985), 319–346.
- RS-2. M. A. RONAN AND S. D. SMITH, Universal presheaves on group geometries, and modular representations, *J. Algebra* **102** (1986), 135–154.

- S-1. S. D. SMITH, Irreducible modules and parabolic subgroups, *J. Algebra* **75** (1985), 286–289.
- S-2. S. D. SMITH, Sheaf homology and complete reducibility, *J. Algebra* **95** (1985), 72–80.
- SV. S. D. SMITH AND H. VÖLKLEIN, A geometric presentation for the adjoint module of  $SL_3(k)$ , *J. Algebra* **127** (1989), 127–138.
- St. R. STEINBERG, “Lectures on Chevalley groups,” Yale University, 1968.
- V-1. H. VÖLKLEIN, On the geometry of the adjoint representation of a Chevalley group, *J. Algebra* **126** (1989), 139–154.
- V-2. H. VÖLKLEIN, The 1-cohomology of the adjoint module of a Chevalley group, *Forum Math.* **1** (1989), 1–13.
- W. W. WONG, Irreducible modular representations of finite Chevalley groups, *J. Algebra* **20** (1972), 355–367.